

Problem Set 9 Solutions

1. (a) Yes, to 0. Applying the weak law of large numbers, we have

$$\mathbf{P}(|U_i - \mu| > \epsilon) \rightarrow 0 \text{ as } i \rightarrow \infty, \text{ for all } \epsilon > 0$$

Here $\mu = 0$ since $X_i \sim U(-1.0, 1.0)$.

- (b) Yes, to 1. Since $W_i \leq 1$, we have for $\epsilon > 0$,

$$\begin{aligned} \lim_{i \rightarrow \infty} \mathbf{P}(|W_i - 1| > \epsilon) &= \lim_{i \rightarrow \infty} \mathbf{P}(\max\{X_1, \dots, X_i\} < 1 - \epsilon) \\ &= \lim_{i \rightarrow \infty} \mathbf{P}(X_1 < 1 - \epsilon) \cdots \mathbf{P}(X_i < 1 - \epsilon) \\ &= \lim_{i \rightarrow \infty} \left(1 - \frac{\epsilon}{2}\right)^i \\ &= 0. \end{aligned}$$

- (c) Yes, to 0.

$$|V_n| \leq \min\{|X_1|, |X_2|, \dots, |X_n|\}$$

but $\min\{|X_1|, |X_2|, \dots, |X_n|\}$ converges to 0 in probability. So, since $|V_n| \geq 0$, $|V_n|$ converges to 0 in probability. To see why $\min\{|X_1|, |X_2|, \dots, |X_n|\}$ converges to 0 in probability, note that:

$$\begin{aligned} \lim_{i \rightarrow \infty} \mathbf{P}(|\min\{|X_1|, \dots, |X_i|\} - 0| > \epsilon) &= \lim_{i \rightarrow \infty} \mathbf{P}(\min\{|X_1|, \dots, |X_i|\} > \epsilon) \\ &= \lim_{i \rightarrow \infty} \mathbf{P}(|X_1| > \epsilon) \cdot \mathbf{P}(|X_2| > \epsilon) \cdots \mathbf{P}(|X_i| > \epsilon) \\ &= \lim_{i \rightarrow \infty} (1 + \epsilon)^i \text{ since } |X_i| \text{ is uniform between 0 and 1} \\ &= 0. \end{aligned}$$

2. Consider a random variable X with PMF

$$p_X(x) = \begin{cases} p, & \text{if } x = \mu - c; \\ p, & \text{if } x = \mu + c; \\ 1 - 2p, & \text{if } x = \mu. \end{cases}$$

The mean of X is μ , and the variance of X is $2pc^2$. To make the variance equal σ^2 , set $p = \frac{\sigma^2}{2c^2}$. For this random variable, we have

$$\mathbf{P}(|X - \mu| \geq c) = 2p = \frac{\sigma^2}{c^2},$$

and therefore the Chebyshev inequality is tight.

3. (a) Let t_i be the expected time until the state HT is reached, starting in state i , i.e., the mean first passage time to reach state HT starting in state i . Note that t_S is the expected number of tosses until first observing heads directly followed by tails. We have,

$$\begin{aligned} t_S &= 1 + \frac{1}{2}t_H + \frac{1}{2}t_T \\ t_T &= 1 + \frac{1}{2}t_H + \frac{1}{2}t_T \\ t_H &= 1 + \frac{1}{2}t_H \end{aligned}$$

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
 Department of Electrical Engineering & Computer Science
6.041/6.431: Probabilistic Systems Analysis
 (Fall 2010)


and by solving these equations, we find that the expected number of tosses until first observing heads directly followed by tails is

$$t_S = 4.$$

- (b) To find the expected number of additional tosses necessary to again observe heads followed by tails, we recognize that this is the mean recurrence time t_{HT}^* of state HT . This can be determined as

$$\begin{aligned}
 t_{HT}^* &= 1 + p_{HT,H}t_H + p_{HT,T}t_T \\
 &= 1 + \frac{1}{2} \cdot 2 + \frac{1}{2} \cdot 4 \\
 &= 4.
 \end{aligned}$$

- (c) Let's consider a Markov chain with states S, H, T, TT , where S is a starting state, H indicates heads on the current toss, T indicates tails on the current toss (without tails on the previous toss), and TT indicates tails over the last two tosses. The transition probabilities for this Markov chain are illustrated below in the state transition diagram:



Let t_i be the expected time until the state TT is reached, starting in state i , i.e., the mean first passage time to reach state TT starting in state i . Note that t_S is the expected number of tosses until first observing tails directly followed by tails. We have,

$$\begin{aligned}
 t_S &= 1 + \frac{1}{2}t_H + \frac{1}{2}t_T \\
 t_T &= 1 + \frac{1}{2}t_H \\
 t_H &= 1 + \frac{1}{2}t_H + \frac{1}{2}t_T
 \end{aligned}$$

and by solving these equations, we find that the expected number of tosses until first observing two consecutive tails is

$$t_S = 6.$$

- (d) To find the expected number of additional tosses necessary to again observe heads followed by tails, we recognize that this is the mean recurrence time t_{TT}^* of state TT . This can be

determined as

$$\begin{aligned} t_{TT}^* &= 1 + p_{TT,HT}t_H + p_{TT,TT}t_{TT} \\ &= 1 + \frac{1}{2} \cdot 6 + \frac{1}{2} \cdot 0 \\ &= 4. \end{aligned}$$

It may be surprising that the average number of tosses until the first two consecutive tails is greater than the average number of tosses until heads is directly followed by tails, considering that the mean recurrence time between pairs of tosses with heads directly followed by tails equals the mean recurrence time between pairs of tosses that are both tails (or equivalently, the long-term frequency of pairs of tosses with heads followed by tails equals the long-term frequency of pairs of tosses with two consecutive tails¹). This is a start-up artifact. Note that the distribution of the first passage time to reach state HT (or TT) starting in state S is the same as the conditional distribution of the recurrence time of state HT (or TT), given that it is greater than 1. Although in both cases the *expected values* of the recurrence times are equal (this is what parts (b) and (d) tell us), the conditional expected values of the recurrence time given that it is greater than 1 is not the same in both cases (possible, because the unconditional distributions are not equal).

4. (a) The long-term frequency of winning can be found as sum of the long-term frequency of transitions from 1 to 2 and 2 to 2. These can be found from the steady-state probabilities π_1 and π_2 , which are known to exist as the chain is aperiodic and recurrent. The local balance and normalization equations are as follows:

$$\begin{aligned} \frac{7}{15}\pi_1 &= \frac{5}{9}\pi_2, \\ \pi_1 + \pi_2 &= 1. \end{aligned}$$

Solving these we obtain,

$$\pi_1 = \frac{25}{46} \approx 0.54, \quad \pi_2 = \frac{21}{46} \approx 0.46.$$

The probability of winning, which is the long-term frequency of the transitions from 1 to 2 and 2 to 2, can now be found as

$$\mathbf{P}(\text{winning}) = \pi_1 p_{12} + \pi_2 p_{22} = \frac{25}{46} \frac{7}{15} + \frac{21}{46} \frac{4}{9} = \frac{21}{46} \approx 0.46.$$

Note that from the balance equation for state 2,

$$\pi_2 = \pi_1 p_{12} + \pi_2 p_{22},$$

the long-term probability of winning always equals π_2 .

- (b) This question is one of determining the probability of absorption into the recurrent class $\{1A, 2A\}$. This probability of absorption can be found by recognizing that it will be the ratio of probabilities

$$\frac{p_{1,1A}}{p_{1,1A} + p_{1,1B}} = \frac{\frac{2}{15}}{\frac{2}{15} + \frac{1}{15}} = \frac{2}{3}.$$

¹See problem 7.34 on page 399 of the text for a detailed explanation of this correspondence between mean recurrence times and steady-state probabilities.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
Department of Electrical Engineering & Computer Science
6.041/6.431: Probabilistic Systems Analysis
(Fall 2010)

More methodically, if we define a_i as the probability of being absorbed into the class $\{1A, 2A\}$, starting in state i , we can solve for the a_i by solving the system of equations

$$\begin{aligned} a_1 &= p_{1,1A} + p_{11}a_1 + p_{12}a_2 \\ &= \frac{2}{15} + \frac{1}{3}a_1 + \frac{7}{15}a_2 \\ a_2 &= p_{21}a_1 + p_{22}a_2 \\ &= \frac{5}{9}a_1 + \frac{4}{9}a_2, \end{aligned}$$

from which we determine that $a_1 = \frac{p_{1,1A}}{p_{1,1A} + p_{1,1B}} = \frac{2}{3}$.

- (c) Let A, B be the events that Jack eventually plays with decks $1A$ & $2A, 1B$ & $2B$, respectively, when starting in state 1. From part (b), we know that $\mathbf{P}(A) = a_1 = \frac{2}{3}$ and $\mathbf{P}(B) = 1 - a_1 = \frac{1}{3}$. The probability of winning can be determined as

$$\mathbf{P}(\text{winning}) = \mathbf{P}(\text{winning}|A)\mathbf{P}(A) + \mathbf{P}(\text{winning}|B)\mathbf{P}(B).$$

By considering the corresponding the appropriate recurrent class and solving a problem similar to part (a), $\mathbf{P}(\text{winning}|A)$ and $\mathbf{P}(\text{winning}|B)$ can be determined; in these cases, the steady-state probabilities of each recurrent class are defined under the assumption of being absorbed into that particular recurrent class. Let's begin with $\mathbf{P}(\text{winning}|A)$. The local balance and normalization equations for the recurrent class $\{1A, 2A\}$ are

$$\begin{aligned} \frac{3}{5}\pi_{1A} &= \frac{1}{5}\pi_{2A}, \\ \pi_{1A} + \pi_{2A} &= 1. \end{aligned}$$

Solving these we obtain,

$$\pi_{1A} = \frac{1}{4}, \quad \pi_{2A} = \frac{3}{4},$$

and hence conclude that

$$\mathbf{P}(\text{winning}|A) = p_{1A,2A}\pi_{1A} + p_{2A,2A}\pi_{2A} = \pi_{2A} = \frac{3}{4}.$$

Similarly, the local balance and normalization equations for the recurrent class $\{1B, 2B\}$ are

$$\begin{aligned} \frac{3}{4}\pi_{1B} &= \frac{1}{8}\pi_{2B}, \\ \pi_{1B} + \pi_{2B} &= 1. \end{aligned}$$

Solving these we obtain,

$$\pi_{1B} = \frac{1}{7}, \quad \pi_{2B} = \frac{6}{7},$$

and hence conclude that

$$\mathbf{P}(\text{winning}|B) = p_{1B,2B}\pi_{1B} + p_{2B,2B}\pi_{2B} = \pi_{2B} = \frac{6}{7}.$$

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
 Department of Electrical Engineering & Computer Science
6.041/6.431: Probabilistic Systems Analysis
 (Fall 2010)

Putting these pieces together, we have that

$$\begin{aligned} \mathbf{P}(\text{winning}) &= \mathbf{P}(\text{winning}|A)\mathbf{P}(A) + \mathbf{P}(\text{winning}|B)\mathbf{P}(B) \\ &= \frac{3}{4} \cdot \frac{2}{3} + \frac{6}{7} \cdot \frac{1}{3} \\ &= \frac{11}{14} \approx 0.79, \end{aligned}$$

meaning that Jack substantially increases the odds to his favor by slipping additional cards into the decks.

- (d) The expected time until Jack slips cards into the deck is the same as the expected time until the Markov chain enters a recurrent state. Let μ_i be the expected amount of time until a recurrent state is reached from state i . We have the equations

$$\begin{aligned} \mu_1 &= 1 + p_{11}\mu_1 + p_{12}\mu_2 = 1 + \frac{1}{3}\mu_1 + \frac{7}{15}\mu_2 \\ \mu_2 &= 1 + p_{21}\mu_1 + p_{22}\mu_2 = 1 + \frac{5}{9}\mu_1 + \frac{4}{9}\mu_2, \end{aligned}$$


which when solved, yields the expected time until Jack slips cards into the deck,

$$\mu_1 = 9.2.$$

- (e) Let S be the number of times that the dealer switches from deck #2 to deck #1, which equals the number of times that he/she switches from deck #1 to deck #2. Let p be the probability that $S = 0$, which is the sum of the probability of all ways for the first change of state to be from state 1 to state 1A or state 1B,

$$p = \left(\frac{2}{15} + \frac{1}{15}\right) + \left(\frac{1}{3}\right) \left(\frac{2}{15} + \frac{1}{15}\right) + \left(\frac{1}{3}\right)^2 \left(\frac{2}{15} + \frac{1}{15}\right) + \dots = \frac{1}{1 - 1/3} \cdot \frac{3}{15} = \frac{3}{10}.$$

Alternatively, p is the probability of absorption of the following modified chain into an absorbing state (1A or 1B), when started in state 1:



As $\mathbf{P}(S > 0) = 1 - p$, and similarly, $\mathbf{P}(S > k + 1 | S > k) = 1 - p$, it should be clear that S will be a shifted geometric, and thus

$$p_S(k) = \left(\frac{7}{10}\right)^k \frac{3}{10} \quad k = 0, 1, 2, \dots$$

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
 Department of Electrical Engineering & Computer Science
6.041/6.431: Probabilistic Systems Analysis
 (Fall 2010)

- (f) Note that S from part (e) is the total number of cycles from 1 to 2 and back to 1. During the i th cycle, the number of wins, W_i , is a geometric random variable with parameter $q = \frac{5}{9}$. Thus the total number of wins by Jack before he slips extra cards into the deck is

$$W = W_1 + W_2 + \dots + W_S,$$

which is a random number of random variables, all of which are independent. Conditioned on $S > 0$, W is a geometric (with parameter p) number of geometric (with parameter q) random variables, all conditionally independent, and thus from the theory of splitting Bernoulli processes,

$$p_{W|S>0}(k) = (1 - pq)^{k-1}pq \quad k = 1, 2, \dots,$$

where $pq = \frac{3}{10} \cdot \frac{5}{9} = \frac{1}{6}$. When $S = 0$, it follows that $W = 0$, and thus by total probability,

$$p_W(k) = \begin{cases} \frac{3}{10} & k = 0 \\ \left(\frac{7}{10}\right)\left(\frac{5}{9}\right)^{k-1}\frac{1}{6} & k = 1, 2, \dots \end{cases}.$$

- (g) Let W be the total number of wins before slipping cards into the deck (as in part (f)), and similarly let L be the total number of losses before absorption. We know from part (d) that $\mathbf{E}[W + L] = \mu_1 = 9.2$. From part (f) we can find $\mathbf{E}[W]$ by total expectation,

$$\mathbf{E}[W] = E[W|S = 0]\mathbf{P}(S = 0) + E[W|S > 0]\mathbf{P}(S > 0) = \frac{7/10}{1/6} = \frac{42}{10} = 4.2,$$

because when conditioned on $S > 0$, the number of wins, W , is a geometric random variable with parameter $pq = \frac{1}{6}$. From linearity of expectation, we find

$$\mathbf{E}[L - W] = \mathbf{E}[W + L] - 2\mathbf{E}[W] = 9.2 - 2 \cdot 4.2 = 0.8.$$

- (h) Using A to again denote the probability of being absorbed into the recurrent class $\{1A, 2A\}$, starting in state 1,

$$\begin{aligned} \mathbf{P}(X_n = 2A|X_{n+1} = 1A) &= \frac{\mathbf{P}(X_{n+1} = 1A|X_n = 2A)\mathbf{P}(X_n = 2A)}{\mathbf{P}(X_{n+1} = 1A)} \\ &= \frac{\mathbf{P}(X_{n+1} = 1A|X_n = 2A)\mathbf{P}(X_n = 2A|A)\mathbf{P}(A)}{\mathbf{P}(X_{n+1} = 1A|A)\mathbf{P}(A)} \\ &\approx \frac{p_{2A,1A}\pi_{2A}}{\pi_{1A}} \\ &= \frac{\frac{1}{5} \cdot \frac{3}{4}}{\frac{1}{4}} \\ &= \frac{3}{5}. \end{aligned}$$

Note that the right hand side above equals $p_{1A,2A}$, as clear from the local balance equation $\pi_{1A}p_{1A,2A} = \pi_{2A}p_{2A,1A}$.

G1[†]. With $a > 0$ and $c \geq 0$,

$$\begin{aligned}\mathbf{P}(X - \mu \geq a) &= \mathbf{P}(X - \mu + c \geq a + c) \\ &\leq \mathbf{P}((X - \mu + c)^2 \geq (a + c)^2) \\ &\leq \frac{\mathbf{E}((X - \mu + c)^2)}{(a + c)^2} \\ &= \frac{(\sigma^2 + c^2)}{(a + c)^2}\end{aligned}$$

where the first inequality follows from the fact that $a + c > 0$, and the second inequality follows from the Markov inequality.

To tighten the bound, we treat $(\sigma^2 + c^2)/(a + c)^2$ as a function of c , and find c such that the derivative is 0. The minimum occurs at $c = \sigma^2/a$. Therefore,

$$\mathbf{P}(X - \mu \geq a) \leq \frac{(\sigma^2 + \frac{\sigma^4}{a^2})}{(a + \frac{\sigma^2}{a})^2} = \frac{\sigma^2}{(\sigma^2 + a^2)}$$

MIT OpenCourseWare
<http://ocw.mit.edu>

6.041SC Probabilistic Systems Analysis and Applied Probability
Fall 2013

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.