

## In-Class Problems Week 11, Fri.

### Problem 1.

We are interested in generating functions for the number of different ways to compose a bag of  $n$  donuts subject to various restrictions. For each of the restrictions in (a)-(e) below, find a closed form for the corresponding generating function.

- (a) All the donuts are chocolate and there are at least 3.
- (b) All the donuts are glazed and there are at most 2.
- (c) All the donuts are coconut and there are exactly 2 or there are none.
- (d) All the donuts are plain and their number is a multiple of 4.
- (e) The donuts must be chocolate, glazed, coconut, or plain and:
  - there must be at least 3 chocolate donuts, and
  - there must be at most 2 glazed, and
  - there must be exactly 0 or 2 coconut, and
  - there must be a multiple of 4 plain.
- (f) Find a closed form for the number of ways to select  $n$  donuts subject to the constraints of the previous part.

### Problem 2. (a) Let

$$S(x) ::= \frac{x^2 + x}{(1 - x)^3}.$$

What is the coefficient of  $x^n$  in the generating function series for  $S(x)$ ?

- (b) Explain why  $S(x)/(1 - x)$  is the generating function for the sums of squares. That is, the coefficient of  $x^n$  in the series for  $S(x)/(1 - x)$  is  $\sum_{k=1}^n k^2$ .
- (c) Use the previous parts to prove that

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}.$$

## Appendix

Let  $[x^n]F(x)$  denote the coefficient of  $x^n$  in the power series for  $F(x)$ . Then,

$$[x^n] \left( \frac{1}{(1 - \alpha x)^k} \right) = \binom{n + k - 1}{k - 1} \alpha^n. \quad (1)$$

### Partial Fractions

Here's a particular case of the Partial Fraction Rule that should be enough to illustrate the general Rule. Let

$$r(x) ::= \frac{p(x)}{(1 - \alpha x)^2(1 - \beta x)(1 - \gamma x)^3}$$

where  $\alpha, \beta, \gamma$  are distinct complex numbers, and  $p(x)$  is a polynomial of degree less than the denominator, namely, less than 6. Then there are unique numbers  $a_1, a_2, b, c_1, c_2, c_3 \in \mathbb{C}$  such that

$$r(x) = \frac{a_1}{1 - \alpha x} + \frac{a_2}{(1 - \alpha x)^2} + \frac{b}{1 - \beta x} + \frac{c_1}{1 - \gamma x} + \frac{c_2}{(1 - \gamma x)^2} + \frac{c_3}{(1 - \gamma x)^3}$$

MIT OpenCourseWare  
<http://ocw.mit.edu>

6.042J / 18.062J Mathematics for Computer Science  
Spring 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.