




Philosophy of QM 24.111

Sixth lecture,
14 Feb. 2005


THE TWO-PATH EXPERIMENT— What we expect:




THE TWO-PATH EXPERIMENT— What we expect:




THE TWO-PATH EXPERIMENT— What we expect:



THE TWO-PATH EXPERIMENT— What we observe:



THE TWO-PATH EXPERIMENT— What we observe:



TWO PROBLEMS:

Our examination of the two-path experiment left us with two **different** problems:

- What is the particle doing when we do not observe which path it follows? Does it somehow follow both paths? Neither path?
- How can we construct a theory that will give us the right prediction?

We will now focus on the *second* problem—
it is much easier than the first!

THE CENTRAL IDEA:

We will use vector spaces to represent

- the physical states of systems of particles;
- the experiments we can perform on these systems.

WARNING: In the literature, these experiments are almost always called “measurements”. Be careful of this word’s connotations!!!

SPIN MEASUREMENTS

First approximation (for the spin state of a **single particle**):


Vector space used to represent states and experiments is **\mathbf{R}^2** .

Unit vectors represent different possible spin states.

Orthogonal axes represent different possible experiments.

One axis corresponds to the “up” outcome and the other to the “down” outcome.

Example:




THE STATISTICAL ALGORITHM

To calculate **Prob(UP)** for 0° experiment:

Project the state-vector onto the “UP, 0° ” axis;

Square the length of this projection.

Answer: $\cos^2(45^\circ) = 1/2$.



To calculate **Prob(DOWN)** for 0° experiment:

Project the state-vector onto the “DOWN, 0° ” axis;

Square the length of this projection.

Answer: $\sin^2(45^\circ) = 1/2$.

Pythagoras' theorem guarantees that these probabilities will sum to 1.

CAUTIOUS INSTRUMENTALISM

We will, for the time being, adopt a cautiously instrumentalist approach to this way of representing physical states. That is, all we take it to “mean”, when we say that the spin state of a particle is represented by such-and-such a vector, is that this vector can be “plugged into” the statistical algorithm so as to yield the correct probabilities for outcomes of spin measurements performed on that particle.

DERIVING THE COS² LAW

If a particle is certain to go up through a magnet with orientation θ_1 , then its probability for going up through a magnet with orientation θ_2 is

$$\cos^2\left(\frac{\theta_1 - \theta_2}{2}\right)$$

DERIVING THE COS² LAW

If $\text{Prob}(\text{UP}, \theta_1) = 1$, then $\text{Prob}(\text{UP}, \theta_2) = f(\theta_1, \theta_2)$.

Assumptions:

1. f depends only on the angle difference $|\theta_1 - \theta_2|$.
2. f is the same with **UP** replaced by **DOWN**.
3. $f(0) = 1$ (of course); $f(\pi) = 0$.
4. f is monotonically decreasing.
5. f is continuous.
6. The values of f are determined in accordance with the statistical algorithm.

FIRST STEP:

If $\text{Prob}(\text{UP}, \theta_1) = 1$, then $\text{Prob}(\text{UP}, \theta_2) = f(\theta_1, \theta_2)$.

Suppose $\text{Prob}(\text{UP}, 0) = 1$.

Then $\text{Prob}(\text{UP}, \pi) = 0$. (by assumptions 1&3)

So $\text{Prob}(\text{DOWN}, \pi) = 1$. (either DOWN or UP must happen)

And $\text{Prob}(\text{UP}, \theta) = f(\theta)$. (by 1)

And $\text{Prob}(\text{DOWN}, \theta) = f(\pi - \theta)$. (by 1 and 2)

But $\text{Prob}(\text{UP}, \theta) + \text{Prob}(\text{DOWN}, \theta) = 1$.

$$\therefore f(\pi - \theta) = 1 - f(\theta).$$


$$\therefore f(\pi/2) = 1/2.$$

1. f depends only on $|\theta_1 - \theta_2|$.
2. f is the same with UP replaced by DOWN.
3. $f(0) = 1$; $f(\pi) = 0$.
4. f is monotonically decreasing.
5. f is continuous.

WHAT THIS SHOWS:

Assumptions 1 - 5 constrain f only this much:

f must be a continuous, monotonically decreasing function with the values $f(0) = 1$, $f(\pi/2) = 1/2$, $f(\pi) = 0$; and $f(\pi - \theta) = 1 - f(\theta)$.




ADDING ASSUMPTION 6:

First, we arbitrarily choose
an axis to represent the
UP,0 outcome...

...and an axis to represent
the **DOWN,0** outcome.


Note that the **DOWN,0**
axis is the same as the
UP, π axis.



ADDING ASSUMPTION 6:


Next, we observe that the $UP, \pi/2$ state-vector must *bisect* these axes:

Notice that the angle it makes to the $UP, 0$ axis is $\pi/4$.




ADDING ASSUMPTION 6:

Next, we observe that the $UP, \pi/4$ axis must be chosen in such a way that the $UP, 0$ and $UP, \pi/2$ state-vectors have projections onto it of *equal length*. There are two such axes:



ADDING ASSUMPTION 6:


But the second choice is ruled out by the requirement that f be monotonically decreasing; for this choice gives us $f(\pi/4) < 1/2$.



ADDING ASSUMPTION 6:

So we have now fixed the $UP,0$, $UP,\pi/4$, $UP,\pi/2$, and UP,π state-vectors (and likewise the UP and $DOWN$ axes for these four directions).

For these directions, the statistical algorithm yields $\text{Prob}(UP,\theta) = \cos^2(\theta/2)$.



ADDING ASSUMPTION 6:

The very same reasoning can now be applied to the directions

$$\theta = 3\pi/4$$


$$\theta = \pi/8$$

$$\theta = 3\pi/8$$

$$\theta = 5\pi/8$$

$$\theta = 7\pi/8$$

...etc. For each $\theta = k\pi/2^n$, the UP, θ state-vector lies at an angle of $\theta/2$ to the $UP, 0$ axis; hence $f(\theta) = \cos^2(\theta/2)$. Since f is continuous, it follows that for every θ , $f(\theta) = \cos^2(\theta/2)$ —i.e., we have derived the \cos^2 -law.




GETTING FANCIER: COMPLEX NUMBERS

It is in fact possible to measure the spin of a particle in *any* direction—not just directions confined to the x-z plane. So there is, for example, a possible spin state in which the particle is certain to go **UP** if spin along the y-direction is measured.

For such a particle, $\text{Prob}(\text{UP},x) = 1/2 = \text{Prob}(\text{UP},z)$. What is its state-vector?

Where can the **UP,y**
state-vector fit?



GETTING FANCIER: COMPLEX NUMBERS

Answer: **It can't.** In order to represent the spin state of such a particle, we need to employ not the vector space \mathbb{R}^2 (the vector space of pairs of real numbers), but the vector space \mathbb{C}^2 (the vector space of pairs of *complex* numbers).

We will mostly ignore this complication
throughout the course.