

24. TRIPLE INTEGRALS

Definition 24.1. Let $B = [a, b] \times [c, d] \times [e, f] \subset \mathbb{R}^3$ be a box in space. A **partition** \mathcal{P} of B is a triple of sequences:

$$\begin{aligned} a &= x_0 < x_1 < \cdots < x_n = b \\ c &= y_0 < y_1 < \cdots < y_n = d \\ e &= z_0 < z_1 < \cdots < z_n = f. \end{aligned}$$

The **mesh** of \mathcal{P} is

$$m(\mathcal{P}) = \max\{x_i - x_{i-1}, y_i - y_{i-1}, z_i - z_{i-1} \mid 1 \leq i \leq n\}.$$

Now suppose we are given a function

$$f: B \longrightarrow \mathbb{R}$$

Pick

$$\vec{c}_{ijk} \in B_{ijk} = [x_{i-1}, x_i] \times [y_{j-1}, y_j] \times [z_{i-1}, z_i].$$

Definition 24.2. The sum

$$S = \sum_{i=1}^n \sum_{j=1}^n \sum_{k=1}^n f(\vec{c}_{ijk})(x_i - x_{i-1})(y_j - y_{j-1})(z_i - z_{i-1}),$$

is called a **Riemann sum**.

Definition 24.3. The function $f: B \longrightarrow \mathbb{R}$ is called **integrable**, with integral I , if for every $\epsilon > 0$, we may find a $\delta > 0$ such that for every mesh \mathcal{P} whose mesh size is less than δ , we have

$$|I - S| < \epsilon,$$

where S is any Riemann sum associated to \mathcal{P} .

If $W \subset \mathbb{R}^3$ is a bounded subset and $f: W \longrightarrow \mathbb{R}$ is a bounded function, then pick a box B containing W and extend f by zero to a function $\tilde{f}: B \longrightarrow \mathbb{R}$,

$$\tilde{f}(x) = \begin{cases} f(x) & \text{if } x \in W \\ 0 & \text{otherwise.} \end{cases}$$

If \tilde{f} is integrable, then we write

$$\iiint_W f(x, y, z) \, dx \, dy \, dz = \iiint_B \tilde{f}(x, y, z) \, dx \, dy \, dz.$$

In particular

$$\text{vol}(W) = \iiint_W 1 \, dx \, dy \, dz.$$

There are two pairs of results, which are much the same as the results for double integrals:

Proposition 24.4. Let $W \subset \mathbb{R}^2$ be a bounded subset and let $f: W \rightarrow \mathbb{R}$ and $g: W \rightarrow \mathbb{R}$ be two integrable functions. Let λ be a scalar.

Then

(1) $f + g$ is integrable over W and

$$\iiint_W f(x, y, z) + g(x, y, z) \, dx \, dy \, dz = \iiint_W f(x, y, z) \, dx \, dy \, dz + \iiint_W g(x, y, z) \, dx \, dy \, dz.$$

(2) λf is integrable over W and

$$\iiint_W \lambda f(x, y, z) \, dx \, dy \, dz = \lambda \iiint_W f(x, y, z) \, dx \, dy \, dz.$$

(3) If $f(x, y, z) \leq g(x, y, z)$ for any $(x, y, z) \in W$, then

$$\iiint_W f(x, y, z) \, dx \, dy \, dz \leq \iiint_W g(x, y, z) \, dx \, dy \, dz.$$

(4) $|f|$ is integrable over W and

$$\left| \iiint_W f(x, y, z) \, dx \, dy \, dz \right| \leq \iiint_W |f(x, y, z)| \, dx \, dy \, dz.$$

Proposition 24.5. Let $W = W_1 \cup W_2 \subset \mathbb{R}^3$ be a bounded subset and let $f: W \rightarrow \mathbb{R}$ be a bounded function.

If f is integrable over W_1 and over W_2 , then f is integrable over W and $W_1 \cap W_2$, and we have

$$\begin{aligned} \iiint_W f(x, y, z) \, dx \, dy \, dz &= \iiint_{W_1} f(x, y, z) \, dx \, dy \, dz + \iiint_{W_2} f(x, y, z) \, dx \, dy \, dz \\ &\quad - \iiint_{W_1 \cap W_2} f(x, y, z) \, dx \, dy \, dz. \end{aligned}$$

Definition 24.6. Define three maps

$$\pi_{ij}: \mathbb{R}^3 \rightarrow \mathbb{R}^2,$$

by projection onto the i th and j th coordinate.

In coordinates, we have

$$\pi_{12}(x, y, z) = (x, y), \quad \pi_{23}(x, y, z) = (y, z), \quad \text{and} \quad \pi_{13}(x, y, z) = (x, z).$$

For example, if we start with a solid pyramid and project onto the xy -plane, the image is a square, but it project onto the xz -plane, the image is a triangle. Similarly onto the yz -plane.

Definition 24.7. A bounded subset $W \subset \mathbb{R}^3$ is an **elementary subset** if it is one of four types:

Type 1: $D = \pi_{12}(W)$ is an elementary region and

$$W = \{ (x, y, z) \in \mathbb{R}^3 \mid (x, y) \in D, \epsilon(x, y) \leq z \leq \phi(x, y) \},$$

where $\epsilon: D \rightarrow \mathbb{R}$ and $\phi: D \rightarrow \mathbb{R}$ are continuous functions.

Type 2: $D = \pi_{23}(W)$ is an elementary region and

$$W = \{ (x, y, z) \in \mathbb{R}^3 \mid (y, z) \in D, \alpha(y, z) \leq x \leq \beta(y, z) \},$$

where $\alpha: D \rightarrow \mathbb{R}$ and $\beta: D \rightarrow \mathbb{R}$ are continuous functions.

Type 3: $D = \pi_{13}(W)$ is an elementary region and

$$W = \{ (x, y, z) \in \mathbb{R}^3 \mid (x, z) \in D, \gamma(x, z) \leq y \leq \delta(x, z) \},$$

where $\gamma: D \rightarrow \mathbb{R}$ and $\delta: D \rightarrow \mathbb{R}$ are continuous functions.

Type 4: W is of type 1, 2 and 3.

The solid pyramid is of type 4.

Theorem 24.8. Let $W \subset \mathbb{R}^3$ be an elementary region and let $f: W \rightarrow \mathbb{R}$ be a continuous function.

Then

(1) If W is of type 1, then

$$\iiint_W f(x, y, z) \, dx \, dy \, dz = \iint_{\pi_{12}(W)} \left(\int_{\epsilon(x, y)}^{\phi(x, y)} f(x, y, z) \, dz \right) \, dx \, dy.$$

(2) If W is of type 2, then

$$\iiint_W f(x, y, z) \, dx \, dy \, dz = \iint_{\pi_{23}(W)} \left(\int_{\alpha(y, z)}^{\beta(y, z)} f(x, y, z) \, dx \right) \, dy \, dz.$$

(3) If W is of type 3, then

$$\iiint_W f(x, y, z) \, dx \, dy \, dz = \iint_{\pi_{13}(W)} \left(\int_{\gamma(x, z)}^{\delta(x, z)} f(x, y, z) \, dy \right) \, dx \, dz.$$

Let's figure out the volume of the solid ellipsoid:

$$W = \left\{ (x, y, z) \in \mathbb{R}^3 \mid \left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 + \left(\frac{z}{c}\right)^2 \leq 1 \right\}.$$

This is an elementary region of type 4.

$$\begin{aligned}
\text{vol}(W) &= \iiint_W dx \, dy \, dz \\
&= \int_{-a}^a \left(\int_{-b\sqrt{1-(\frac{x}{a})^2}}^{b\sqrt{1-(\frac{x}{a})^2}} \left(\int_{-c\sqrt{1-(\frac{x}{a})^2-(\frac{y}{b})^2}}^{c\sqrt{1-(\frac{x}{a})^2-(\frac{y}{b})^2}} dz \right) dy \right) dx \\
&= \int_{-a}^a \left(\int_{-b\sqrt{1-(\frac{x}{a})^2}}^{b\sqrt{1-(\frac{x}{a})^2}} 2c\sqrt{1-(\frac{x}{a})^2-(\frac{y}{b})^2} dy \right) dx \\
&= 2c \int_{-a}^a \left(\int_{-b\sqrt{1-(\frac{x}{a})^2}}^{b\sqrt{1-(\frac{x}{a})^2}} \sqrt{1-(\frac{x}{a})^2-(\frac{y}{b})^2} dy \right) dx \\
&= \frac{2c}{b} \int_{-a}^a \left(\int_{-b\sqrt{1-(\frac{x}{a})^2}}^{b\sqrt{1-(\frac{x}{a})^2}} \sqrt{b^2 \left(1-(\frac{x}{a})^2\right) - y^2} dy \right) dx \\
&= \frac{\pi c}{b} \int_{-a}^a b^2 \left(1-(\frac{x}{a})^2\right) dx \\
&= \pi bc \int_{-a}^a 1-(\frac{x}{a})^2 dx \\
&= \pi bc \left[x - \frac{x^3}{3a^2} \right]_{-a}^a \\
&= \pi bc \left(2a - 2\frac{a^3}{3a^2} \right) \\
&= \frac{4\pi}{3} abc.
\end{aligned}$$

MIT OpenCourseWare
<http://ocw.mit.edu>

18.022 Calculus of Several Variables
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.