

18.445 Introduction to Stochastic Processes

Lecture 10: Hitting times

Hao Wu

MIT

16 March 2015

Recall

- Consider a network $(G = (V, E), \{c(e) : e \in E\})$. The effective resistance is defined by

$$R(a \leftrightarrow z) = (W(a) - W(z)) / \|I\|.$$

- Consider a random walk on the network, the Green's function is defined by

$$G_{\tau}(a, x) = \mathbb{E}[\#\text{visits to } x \text{ before } \tau].$$

- We have that

$$G_{\tau_z}(a, a) = c(a)R(a \leftrightarrow z).$$

Today's Goal

- hitting time
- commute time
- transitive network

Target time

Suppose that $(X_n)_{n \geq 0}$ is an irreducible Markov chain with transition matrix P and stationary measure π . Let τ_x be the hitting time :

$$\tau_x = \min\{n \geq 0 : X_n = x\}.$$

Lemma

The quantity

$$\sum_x \mathbb{E}_a[\tau_x] \pi(x)$$

does not depend on a ; and we call it target time and denote it by t_\odot .

Hitting time

Definition

$$t_{hit} := \max_{x,y} \mathbb{E}_x[\tau_y] \geq t_{\odot}.$$

Lemma

Suppose that the chain is irreducible with stationary measure π . Then

$$t_{hit} \leq 2 \max_w \mathbb{E}_{\pi}[\tau_w].$$

Theorem

For an irreducible transitive Markov chain, we have

$$t_{hit} \leq 2t_{\odot}.$$

Transitive Markov chain

Roughly, a transitive Markov chain “looks the same” from any point in the state space.

Definition

A Markov chain is called transitive if for each pair $(x, y) \in \Omega \times \Omega$, there is a bijection $\varphi : \Omega \rightarrow \Omega$ such that

$$\varphi(x) = y; \quad P(\varphi(z), \varphi(w)) = P(z, w), \forall z, w.$$

Example : simple random walk on N -cycle, on hypercube.

Lemma

For a transitive Markov chain on finite state space Ω , the uniform measure is stationary.

Commute time

Definition

Suppose that the Markov chain starts from $X_0 = a$. The commute time between a and b is defined by

$$\tau_{ba} = \min\{n \geq \tau_b : X_n = a\}.$$

Theorem (Commute Time Identity)

Consider a random walk on the network $(G = (V, E), \{c(e) : e \in E\})$, we have

$$\mathbb{E}_a[\tau_{ba}] = \mathbb{E}_a[\tau_b] + \mathbb{E}_b[\tau_a] = c_G R(a \leftrightarrow b).$$

Lemma

Suppose that the Markov chain is irreducible with stationary measure π . Suppose that τ is a stopping time satisfying $\mathbb{P}_a[X_\tau = a] = 1$. Then

$$G_\tau(a, x) = \mathbb{E}_a[\tau]\pi(x).$$

Transitive network

Generally, $\mathbb{E}_a[\tau_b]$ and $\mathbb{E}_b[\tau_a]$ can be very different (see Exercise 10.3). However, if the network is transitive, they are equal.

Definition

A network $(G = (V, E), \{c(e) : e \in E\})$ is transitive if for each pair $(x, y) \in V \times V$, there exists a bijection $\varphi : V \rightarrow V$ such that

$$\varphi(x) = y; \quad c(\varphi(z), \varphi(w)) = c(z, w), \forall z, w.$$

Remark : The random walk on a transitive network is a transitive Markov chain.

Theorem

For the random walk on a transitive (connected) network, for any vertices a and b , we have

$$\mathbb{E}_a[\tau_b] = \mathbb{E}_b[\tau_a].$$

Summary

For random walk on network

- $t_{\odot} \leq t_{hit} \leq 2 \max_w \mathbb{E}_{\pi}[\tau_w]$.
- $\mathbb{E}_a[\tau_{ba}] = c_G R(a \leftrightarrow b)$.

For random walk on transitive network

- $t_{\odot} \leq t_{hit} \leq 2t_{\odot}$.
- $\mathbb{E}_a[\tau_b] = \mathbb{E}_b[\tau_a]$.
- $2\mathbb{E}_a[\tau_b] = c_G R(a \leftrightarrow b)$.

MIT OpenCourseWare
<http://ocw.mit.edu>

18.445 Introduction to Stochastic Processes

Spring 2015

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.