

MIT OpenCourseWare
<http://ocw.mit.edu>

18.726 Algebraic Geometry
Spring 2009

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

18.726: Algebraic Geometry (K.S. Kedlaya, MIT, Spring 2009)
Higher Riemann-Roch

In this lecture, we discuss some higher-dimensional versions of the Riemann-Roch theorem: the Riemann-Roch theorem for surfaces, the Hirzebruch-Riemann-Roch theorem, and the Grothendieck-Riemann-Roch theorem. For the first, see Hartshorne V.1; for the others, see Chapter 15 of Fulton's *Intersection Theory* (a well-deserved winner of the Steele Prize for mathematical exposition).

1 Surfaces

Let X be a smooth irreducible projective surface over an algebraically closed field k . Let K be a canonical divisor on X . As in the case of curves, the Riemann-Roch theorem combines an input from Serre duality with an Euler characteristic calculation.

The input from Serre duality is that for any divisor D ,

$$H^0(X, \mathcal{L}(D)^\vee \otimes \omega_X) \cong H^2(X, \mathcal{L}(D))^\vee.$$

We can thus write the Euler characteristic $\chi(X, \mathcal{L}(D))$ as

$$\dim_k H^0(X, \mathcal{L}(D)) - \dim_k H^1(X, \mathcal{L}(D)) + \dim_k H^0(X, \mathcal{L}(K - D)).$$

Unfortunately, we can't do much with the term $\dim_k H^1(X, \mathcal{L}(D))$ other than give it a name: it's called the *superabundance* of D . However, we do at least know that it is nonnegative, and this turns out to be surprisingly useful.

The Euler characteristic calculation is made as follows. Write D as the difference between two effective divisors $C - E$ with no common components. We then have exact sequences

$$0 \rightarrow \mathcal{L}(C - E) \rightarrow \mathcal{L}(C) \rightarrow \mathcal{L}(C) \otimes \mathcal{O}_E \rightarrow 0, \quad 0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{L}(C) \rightarrow \mathcal{L}(C) \otimes \mathcal{O}_C \rightarrow 0.$$

By additivity of χ , we get

$$\chi(X, \mathcal{L}(C - E)) = \chi(X, \mathcal{O}_X) + \chi(C, \mathcal{L}(C)) - \chi(E, \mathcal{L}(C)).$$

The first term we are happy to leave alone since it depends only on X . The other two are calculated using *intersection theory* on the surface X . For instance, the term $\chi(E, \mathcal{L}(C))$ equals $C \cdot E + 1 - g_E$, where g_E is the genus and $C \cdot E$ is the length of the scheme-theoretic intersection $C \times_X E$ (this amounts to Riemann-Roch on the curve E).

The term $\chi(C, \mathcal{L}(C))$ is a bit trickier: it is $C \cdot C + 1 - G_C$ where $C \cdot C = C^2$ is the *self-intersection* of C . That can be defined as $C \cdot C'$ if C is linearly equivalent to a divisor C' having no common components with C , but that is not always possible. In fact, the correct definition is to force the intersection pairing to be bilinear, and this sometimes involves letting C^2 take negative values. For instance, if you blow up P^2 at a point, the exceptional divisor has self-intersection -1 . (This is a general pattern; one can in fact blow down any curve isomorphic to \mathbb{P}^1 with self-intersection -1 .)

Moreover, one can write the genera of C and E in terms of the canonical divisor K , basically using Riemann-Roch again:

$$C \cdot (C + K) = 2g_C - 2, \quad E \cdot (E + K) = 2g_E - 2.$$

So

$$\chi(X, \mathcal{L}(D)) = \frac{1}{2}D \cdot (D - K) + \chi(X, \mathcal{O}_X).$$

As in the case of curves, this is useful for many calculations involving the geometry of surfaces, such as the Hodge index theorem and the Nakai-Moishezon criterion. These in turn figure in the classification of surfaces (which you should read about in Hartshorne if you are interested in Abhinav's work).

Theorem (Hodge index theorem). *Fix a projective embedding of X , and let H be a divisor with $\mathcal{L}(H) \cong \mathcal{O}_X(1)$. Then for any divisor D such that $D \cdot H = 0$, we have $D^2 \leq 0$. (This also holds if H is ample, i.e., some positive multiple of H comes from an $\mathcal{O}_X(1)$.)*

Theorem (Nakai-Moishezon criterion). *A divisor D on X is ample if and only if $D^2 > 0$ and $D \cdot C > 0$ for all irreducible curves C on X .*

2 Hirzebruch's generalization

Hirzebruch noticed that the Euler characteristic aspect of Riemann-Roch could be generalized to handle arbitrary vector bundles on arbitrary smooth varieties over an algebraically closed field k . Let me state his result and then explain what it means.

Theorem (Hirzebruch). *Let X be a smooth proper scheme over k . Let \mathcal{F} be a locally free coherent sheaf on X . Then*

$$\chi(X, \mathcal{F}) = \int_X \text{ch}(\mathcal{F}) \cdot \text{td}(T_X).$$

Here T_X is the *tangent bundle* of X , i.e., the dual to the bundle ω_X of Kähler differentials (which is also called the *cotangent bundle*).

The *Chern character* ch is a certain map from coherent sheaves on X to a certain group of *cycles* on X . The latter are formal \mathbb{Q} -linear combinations of subschemes of X modulo a relation of *rational equivalence*. You should imagine this as generalizing the function taking a line bundle \mathcal{L} on a curve C to (the equivalence class of) the divisor of a nonzero rational section of \mathcal{L} .

The group of cycles is graded by codimension, and forms a commutative ring under the (appropriately defined) intersection pairing with the identity being the class of X itself in codimension 0. The Chern character is usually split up as $\sum_d c_d(\cdot)$ with c_d being the bit in codimension d ; for \mathcal{F} locally free of rank 1, we always have

$$c_d(\mathcal{F}) = \frac{1}{d!} c_1(\mathcal{F})^d.$$

The *Todd class* td is another such map on coherent sheaves, which I won't try to construct here, except to give the characterizing identity: for \mathcal{F} locally free of rank d ,

$$\text{td}(\mathcal{F}) \cdot \sum_{p=0}^d (-1)^p \text{ch}(\wedge^p \mathcal{F}^\vee) = c_d(\mathcal{F}).$$

. The point is that it depends only on X , not on \mathcal{F} .

The Chern character and the Todd class are both examples of *characteristic classes* of vector bundles, which originally appeared in algebraic topology as tools for classifying manifolds. For instance, Milnor uses them to construct differentiable manifolds which are homeomorphic but not diffeomorphic to the 7-sphere, the so-called *exotic 7-spheres*. See Milnor and Stasheff, *Characteristic Classes* for an introduction.

Oh, and \int_X means use intersection theory (which is a pretty complicated thing to define, as evidenced by the length of Fulton's book), keep only the zero-dimensional part, and count points.

3 Grothendieck's generalization

In characteristic fashion, Grothendieck noticed that one can make a relative version of the Hirzebruch-Riemann-Roch theorem. Also, one can drop the locally free condition.

Theorem (Grothendieck). *Let $f : X \rightarrow Y$ be a proper morphism of smooth schemes over an algebraically closed field k . Then for any coherent sheaf \mathcal{F} on X ,*

$$\text{ch}(f_*\mathcal{F}) \cdot \text{td}(T_Y) = f_*(\text{ch}(\mathcal{F}) \cdot \text{td}(T_X)).$$

One has to define direct image for cycles; I won't try here.

It should be noted that already our formulation of Hirzebruch's statement is Grothendieck's; the original statement was made in the language of topology. One byproduct of this work is the development of *K-theory*, which is now a frequently occurring construction in both algebraic topology and algebraic geometry.