

18.905 Problem Set 4

Due Wednesday, October 4 in class

1. Let A be the 1-skeleton of Δ^3 , i.e. the union of the vertices and lines. Let CA , the cone on A , be the union of all the lines joining A to the center of Δ^3 . Compute the homology groups $H_*(CA, A)$.
2. Hatcher, exercise 17 on page 132.
3. Hatcher, exercise 27 on page 133.
4. A *generalized homology theory* E is a collection of functors E_n from the category of pairs (X, A) to the category of abelian groups, together with natural transformations $\partial_E : E_n(X, A) \rightarrow E_{n-1}(A, \emptyset)$ which satisfy the Eilenberg-Steenrod axioms except possibly for the dimension axiom. Homology H is one example. In the previous problem set we showed that H^Y , given by functors $H_n^Y(X, A) = H_n(X \times Y, A \times Y)$, is a generalized homology theory.

A *map* of generalized homology theories $\phi : E \rightarrow F$ is a collection of natural transformations $\phi_n : E_n \rightarrow F_n$ such that $\partial_F \circ \phi_n = \phi_{n-1} \circ \partial_E$.

Show that, with this definition, generalized homology theories form a category. Given a map $f : Y \rightarrow Z$ of spaces, explain why we get a map of generalized homology theories $H^f : H^Y \rightarrow H^Z$. Show that this is a functor from spaces to generalized homology theories.