


18.01 Practice Exam 4 solutions

$$\boxed{1} \quad \int_0^{1/2} \frac{x^2}{\sqrt{1-x^2}} dx = \int_0^{\pi/6} \frac{\sin^2 u}{\cos u} \cos u du$$

Put $x = \sin u$ $\frac{1}{2} = \sin \frac{\pi}{6}$
 (or $x = \cos u$) 

$$= \int_0^{\pi/6} \frac{1 - \cos 2u}{2} du$$

$$= \left[\frac{u}{2} - \frac{\sin 2u}{4} \right]_0^{\pi/6} = \frac{\pi}{12} - \frac{\sqrt{3}}{8}$$

$$\boxed{2} \quad y = e^x$$


$$\text{Volume} = \int_0^1 2\pi x e^x dx$$

$$\int x e^x dx = x e^x - \int e^x dx = x e^x - e^x$$

$$\therefore \text{volume} = 2\pi (x e^x - e^x) \Big|_0^1 = 2\pi (0 - (-1)) = 2\pi$$

$$\boxed{3} \quad \frac{4x}{(x^2-1)(x-1)} = \frac{4x}{(x-1)^2(x+1)}$$

$$= \frac{2}{(x-1)^2} + \frac{B+1}{x-1} + \frac{-1}{x+1}$$

by coverup by coverup

Put $x=0$: $0 = 2 - B - 1$; $B=1$

Integrating:

$$\int \frac{4x dx}{(x^2-1)(x+1)} = \frac{-2}{x-1} + \ln(x-1) - \ln(x+1) + C$$

$$= \frac{-2}{x-1} + \ln\left(\frac{x-1}{x+1}\right) + C$$

$$\boxed{4} \quad \int_a^b \sqrt{1+y^2} dx \quad y = \sin x$$

$$y' = 2 \sin x \cos x = \sin 2x$$


Arclength

$$= \int_0^{\pi/4} \sqrt{1 + \sin^2 2x} dx$$

Since $0 \leq \sin^2 2x \leq 1$ on the interval,


$$\int_0^{\pi/4} \sqrt{1 + \sin^2 2x} dx \leq \frac{\pi}{4} \cdot \sqrt{2} < \frac{3.2 \cdot 3}{4 \cdot 2}$$

length of interval 1.2

$$\boxed{5} \quad$$



a) $(x-b)^2 + y^2 = a^2$ (by translation of $x^2 + y^2 = a^2$ to $(b, 0)$ center)
 $x^2 + y^2 - 2bx + b^2 = a^2$

$$\boxed{r^2 - 2br \cos \theta = a^2 - b^2}$$

b) Applying law of cosines to  we get

$$a^2 = r^2 + b^2 - 2br \cos \theta$$

same as part (a).


$$\boxed{6} \quad$$


shows this is a right Δ , $\alpha = \pi/4$
 (or you can solve $\begin{cases} r = 2 \cos \theta \\ r = \sqrt{2} \end{cases}$ simultaneously, for θ)

Using symmetry: $\pi/4$

$$\text{Area} = 2 \cdot \frac{1}{2} \int_0^{\pi/4} (2 \cos \theta)^2 - (\sqrt{2})^2 d\theta$$

$$b) = 2 \int_0^{\pi/4} (2 \cos^2 \theta - 1) d\theta = 2 \cdot \frac{\sin 2x}{2} \Big|_0^{\pi/4} = 1$$

[By elem. geometry:  $= \frac{1}{2} + \frac{\pi}{4}$
 $= \frac{\pi(\sqrt{2})^2}{8} + A \therefore A = \frac{1}{2}, 2A = 1$